

PEDIATRIC SURGERY Update © Vol. 22 No. 02 FEBRUARY 2004

Forme Fruste Choledochal Cyst

In 1985, a new variant of choledochal cyst known as forme fruste was described in the pediatric literature. Forme fruste choledochal cyst (FFCC) is characterized by minimal dilatation of the extrahepatic bile duct which does not grows with time. The normal diameter of the common bile duct in children ranges between two and 6 mm. FFCC is associated with a diameter above six mm and below 10 mm. Most patients with FFCC have a long common channel, in which the common bile duct-pancreatic duct junction is away from the duodenal papilla, with partial obstruction of the terminal common bile duct. FFCC is associated with fever, jaundice, abdominal pain, recurrent pancreatitis and altered liver function tests. Histologically FFCC demonstrates thickened fibrous connective tissue, absent muscular layer with flattened, ulcerated and dysplastic mucosa. Diagnosis is established with ultrasound. ERCP or better yet MRCP can help delineate the anatomy and presence of a long common pancreaticobiliary channel in FFCC. Management consists of cyst excision and Roux-en-y hepaticojejunostomy. Due to the small size the anastomosis is technically difficult and should be performed carefully to avoid stricture and postoperative cholangitis. To maintain ductal anastomosis patency it is imperative that diseased ductal tissue not be incorporated in the anastomosis, the circumstance most likely responsible for the high incidence of anastomotic stricture in choledochal cyst past drainage operations.

References:

1- Lilly JR, Stellin GP, Karrer FM: Forme fruste choledochal cyst. J Pediatr Surg 20(4):449-51, 1985

2- Okada A: Forme fruste choledochal cyst. J Pediatr Surg 21(4):383, 1986

3- Okada A, Oguchi Y, Kamata S, et al: Common channel syndrome - diagnosis with endoscopic retrograde cholangio-pancreatography and surgical treatment. Surgery 93: 634-642, 1983

4- Ando H, Ito T, Nagaya M, Watanabe Y, Seo T, Kaneko K: Pancreaticobiliary Maljunction without choledochal cysts in Infants and Children: Clinical features and surgical therapy. J Pediatr Surg 30(12): 1658-1662, 1995 5- Thomas S, Sen S, Zachariah N, Chacko J, Thomas G: Choledochal cyst sans cyst--experience with six "forme fruste" cases. Pediatr Surg Int 18(4):247-51, 2002

6- Shimotakahara A, Yamataka A, Kobayashi H, Okada Y, Yanai T, Lane GJ, Miyano T: Forme fruste choledochal cyst: long-term follow-up with special reference to surgical technique. J Pediatr Surg 38(12):1833-6, 2003

Gallbladder Polyps

A polypoid lesion identified in the gallbladder of a child is a very rare event. It represents an elevated lesion of the mucosal surface of the gallbladder which in most instances causes parental concern. Fortunately, most polypoid lesions identified in gallbladders are benign (90%). Histologically they are either adenomatous, hyperplastic, gastric heterotopia or cholesterol polyps. The prevalence of such polyps is greater among males and obese children. Ultrasonography is the image method of choice in diagnosing gallbladders polyps in children and adults. They are seen as pedunculated or sessile echogenic lesions attached to the gallbladder wall protruding toward the lumen and fixed in changed of posture. Gallbladder polyps can be associated with acalculous cholecystitis. Lesions smaller than 10 mm do not progress to malignancy or development of stones, and none produces symptoms or complications of biliary disease. Surgical management of gallbladder polyps is indicated when the size of the polypoid lesion is above 10 mm in diameter, when associated with gallstones and when the child has consistent biliary symptoms. Treatment consists of laparoscopic cholecystectomy. Asymptomatic small polyps (< 10 mm) should be maintained under ultrasonographic surveillance.

References:

1- Yang HL, Sun YG, Wang Z: Polypoid lesions of the gallbladder: diagnosis and indications for surgery. Br J Surg 79(3):227-9, 1992

2- Barzilai M, Lerner A: Gallbladder polyps in children: a rare condition. Pediatr Radiol 27(1):54-6, 1997

3- Stringel G, Beneck D, Bostwick HE: Polypoid lesions of the gallbladder in children. JSLS 1(3):247-9, 1997 4- Csendes A, Burgos AM, Csendes P, Smok G, Rojas J: Late follow-up of polypoid lesions of the gallbladder smaller than 10 mm. Ann Surg 234(5):657-60, 2001

5- Kikiros C, Arunachalam P, Lam MH: Adenomatous hyperplastic polyp of the gall bladder associated with cholelithiasis in a child. Pediatr Surg Int 19(1-2):118-9, 2003

6- Stringer MD, Ceylan H, Ward K, Wyatt JI: Gallbladder Polyps in Children - Classification and Management. J Pediatr Surg 38(11): 1680-1684, 2003

Gastrocutaneous Fistula

Gastrocutaneous fistula (GCF) is most commonly identified after removing long standing gastrostomy tubes in children. Other times is the result of gastrojejunal tubes and Crohn's disease. After removing a temporary gastrostomy tube most stomas close between three and six weeks after removal. Persistence of stomach leakage through the gastrostoma is a nuisance, erodes the surrounding skin and causes nutritional depletion. GCF does not close spontaneously when the stoma has been used for a long period of time, when there is distal obstruction (delayed gastric emptying), foreign body reaction (silk), epithelization of the tract (multiple granulomas formation), or associated chronic granulomatous disease (Crohn). Silk suture should be avoided when constructing surgical gastrostomies. When the tube is in place for more than nine months before removal the incidence of GCF can be as high as 45%. Initial non-surgical therapy should include H2-antagonist therapy and silver nitrate cauterization. If this does not work permanent management of GCF consists of surgical closure.

References:

1- Kobak GE, McClenathan DT, Schurman SJ: Complications of removing percutaneous endoscopic gastrostomy tubes in children. J Pediatr Gastroenterol Nutr 30(4):404-7, 2000

2- Gordon JM, Langer JC: Gastrocutaneous fistula in children after removal of gastrostomy tube: incidence and predictive factors. J Pediatr Surg 34(9):1345-6, 1999

3- Aronian JM, Redo SF: Gastrocutaneous fistula after tube gastrostomy. Incidence in infants and children. N Y State J Med 74(13):2364-6, 1974

4- Davies BW, Watson AR, Coleman JE, Rance CH: Do gastrostomies close spontaneously? A review of the fate of gastrostomies following successful renal transplantation in children. Pediatr Surg Int 17(4):326-8, 2001

* Edited by: Humberto Lugo-Vicente, MD, FACS, FAAP Professor /Academic Director of Pediatric Surgery, University of Puerto Rico - School of Medicine, Rio Piedras, Puerto Rico. Address: P.O. Box 10426, Caparra Heights Station, San Juan, Puerto Rico USA 00922-0426. Tel (787)-786-3495 Fax (787)-720-6103 E-mail: *titolugo@coqui.net* Internet: http://home.coqui.net/titolugo

© PSU 1993-2004 ISSN 1089-7739